

# Search for Pentaquarks at CLAS in Photoproduction from Proton

*M. Battaglieri, R. De Vita, V. Kubarovsky, D. Weygand  
and the CLAS Collaboration*



April Meeting




# Quarks and QCD

quarks are among the few known elementary particles

quarks interact by the strong force through the exchange of gluons forming hadrons, the building blocks of ordinary matter

known quark configurations are baryons ( $3q$ ) and mesons ( $q\bar{q}$ )

Quantum Chromo Dynamics allows hadronic states with different quark configuration ( $4q$ ,  $5q$ ,  $3qg$ )



# Exotics and Pentaquarks

**Pentaquarks** are hadronic states with minimal quark content of 5 quarks


$$4q + \bar{q}$$

When the antiquark has a different flavor than the other 4 quark, the pentaquark has “**exotic**” quantum numbers

**Example:**  $uudd\bar{s}$

$$\text{Strangeness} = 0 + 0 + 0 + 0 + 1 = +1$$

An antidecuplet of pentaquarks with masses of 1.5–2.1 GeV was predicted by D. Diakonov et al.



# Evidence for Pentaquark States

- ◆ First evidence for a possible  $\Theta^+(1540)$  was reported by the LEPs/Spring-8 Collaboration in photoproduction off nuclei
- ◆ Corroborated by several experiments ( CLAS, COSY-TOF, DIANA, HERMES, JINR, NOMAD,  $\nu$ BC, SAPHIR , SVD-2, ZEUS) using different probes and targets
- ◆ Positive results have limited event samples in the observed structures
- ◆ Comparison of results from different experiments shows discrepancies in the observed masses
- ◆ Null results have been recently presented (ALEPH, BABAR, BELLE, BES, CDF, COMPASS, DELPHI, E690, FOCUS, HERA-B, PHENIX, SPHINX)

**Definitive confirmation from high statistics/ high resolution experiments is needed to resolve the controversy about the existence of the  $\Theta^+(1540)$  pentaquark**

# Search for Pentaquarks at CLAS

A comprehensive program to search for pentaquarks with high statistics and high resolution **photoproduction experiments** is in progress at **Jefferson Lab**

New experiments seeking evidence of pentaquarks with the **CLAS** detector were approved in 2003–2004 with the goal of confirming previous results and explore new kinematics with at least **a factor 10 increase in statistics**

**g10**      **deuteron**       $E_\gamma \sim 1.0\text{--}3.5 \text{ GeV}$   
data taking completed in 2004


**g11**      **proton**       $E_\gamma \sim 1.6\text{--}3.8 \text{ GeV}$   
data taking completed in 2004

**eg3**      **deuteron**       $E_\gamma \sim 4.0\text{--}5.4 \text{ GeV}$   
data taking completed in 2005


**Super-g**      **proton**       $E_\gamma \sim 3.8 - 5.7 \text{ GeV}$   
planned for 2006

# $\Theta^+$ Search in Exclusive Photoproduction off the proton

J. Barth et al. Phys Lett. B572 2003 127




$$\gamma p \rightarrow \Theta^+ \bar{K}^0$$



$$E_\gamma = 1.6 - 2.6 \text{ GeV}$$


V. Kubarovsky et al. PRL 92 032001 (2004)



$$\gamma p \rightarrow \Theta^+ K^- \pi^+$$


to be tested by a dedicated  
experiment (Super-g)  
planned for 2006

# Jefferson Lab




$E_{\text{max}}$	$\sim 6 \text{ GeV}$
$I_{\text{max}}$	$\sim 200 \mu\text{A}$
Duty Factor	$\sim 100\%$
$\sigma_E/E$	$\sim 2.5 \cdot 10^{-5}$
Beam P	$\sim 80\%$
$E_\gamma$	$\sim 0.8 - 5.7 \text{ GeV}$

CLAS







# The CEBAF Large Acceptance Spectrometer CLAS



## Performance

- ◆  $L = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
- ◆  $\int B \, dl = 2.5 \text{ T m}$
- ◆  $\Delta p/p \sim 0.5\text{--}1 \%$
- ◆  $\sim 4\pi$  acceptance
- ◆ Best suited for multiparticle final states
- ◆ Bremsstrahlung Photon Tagger ( $\Delta E_\gamma/E_\gamma \sim 10^{-3}$ )





# g11@JLab: Spectroscopy of Exotic Baryons with CLAS

## Search for Ground and Excited States


### Proposed measurement and Primary Goals:

- ▶ search for  $\Theta^+(1540)$  and possible excited states in  $\gamma$ -p interaction above threshold ( $E_\gamma = 1.6 - 3.8$  GeV)
- ▶ collect more than 10 times the statistics of previous measurements in the same kinematics
- ▶ establish the mass of any observed peak with 2 MeV accuracy
- ▶ determine total and differential cross section

### Status of the experiment

- ▶ New experiment approved by JLab PAC25 in January 2004.
- ▶ Run in May–July 2004, with a total of  $7 \cdot 10^9$  triggers recorded (Luminosity  $\sim 70$  pb $^{-1}$ )
- ▶ Data calibration and processing completed in January 2005
- ▶ Preliminary results for this reaction

$$\gamma p \rightarrow \Theta^+ \bar{K}^0$$




$$\Theta^+ \rightarrow n K^+$$



$$\bar{K}^0 \rightarrow K_S \rightarrow \pi^+ \pi^-$$

# g11 data quality

Particle  
identification









Reaction  
identification  
and  
kinematical  
coverage






$$\gamma p \rightarrow \pi^+ \pi^- K^+ n$$

*1% of statistics*




# Cross Section Extraction



- Different final states are measured simultaneously in the CLAS detector
- Cross section for known reactions have been extracted to test the accuracy of the analysis procedure
- 1% of statistics






# Search for the $\Theta^+$ (1540)



- ▶ the  $K^0$  is detected via its  $K_S$  component decaying into  $\pi^+ \pi^-$
- ▶ final state is identified using the missing mass technique
- ▶ strangeness is tagged detecting the  $K^+$
- ▶ using the full statistics ( $70 \text{ pb}^{-1}$ ) a total of  $\sim 120000$  events are selected



# Hyperon Production




$$N_{\Lambda} = 91300$$

$$N_{\Sigma^+} = 36200$$

$$N_{\Sigma^-} = 26800$$






# $nK^+$ Mass Spectrum




- ▶ the  $nK^+$  mass spectrum is smooth
- ▶ no structure is observed at a mass of  $\sim 1540$  MeV

# $nK^+$ Mass Spectrum






- ♦ mass vs. CM angle distribution shows coverage of the whole kinematics
- ♦ no evidence for a structure is found for different CM angle ranges



# Evaluation of an Upper Limit on the $\Theta^+$ Cross section


- ◆  $\Theta^+$  signal is searched for as narrow resonance over a smooth background in the  $(nK^+)$  spectrum
- ◆ Resonance width inferred from MC simulations, assuming a negligible intrinsic width
- ◆ Background described as 5<sup>th</sup> order polynomial
- ◆ Signal and background yields extracted fitting:
  - ◇ binned/unbinned spectra
  - ◇ including/excluding  $\Theta^+$  mass region
- ◆  $M(nK^+)$  range (1520 – 1600 MeV) scanned in 5 MeV steps
- ◆ Upper limit derived using Feldman and Cousins approach

# Evaluation of an Upper Limit on the $\Theta^+$ Yield



# $\Theta^+(1540)/\Lambda^*$ Yields


$$\frac{N(\Theta^+)}{N(\Lambda^*)} < 0.2\%$$



# Upper Limit on the $\Theta^+(1540)$ Cross section *energy and angular dependence*


$$\sigma(95\%CL) \propto \frac{\text{Yield}(95\%CL)}{\text{efficiency} \cdot \text{luminosity} \cdot \text{BR}(\Theta^+ \rightarrow nK^+)}$$

- ▶ Absolute cross section measurement
- ▶  $\text{BR}(\Theta^+ \rightarrow nK^+) = 50\%$
- ▶ CLAS detection efficiency evaluated using different  $\Theta^+$  production models





# Upper Limit on the $\Theta^+$ Cross section *mass dependence*



Upper limit (95% CL)

$$\sigma_{\gamma p \rightarrow \Theta^+ K^0} < 1-4 \text{ nb}$$

# Comparison with SAPHIR results

## Observed Yields


### SAPHIR

$$N(\Theta^+)/N(\Lambda^*) \sim 9\%$$

### CLAS

$$N(\Theta^+)/N(\Lambda^*) < 0.5\% \quad (95\% \text{CL})$$

## SAPHIR



## Cross Sections

### SAPHIR


$$\sigma_{\gamma p \rightarrow \Theta^+ K^0} \sim 300 \text{ nb}$$

reanalysis 50 nb

### CLAS

$$\sigma_{\gamma p \rightarrow \Theta^+ K^0} < 1-4 \text{ nb}$$

## g11@CLAS



# Summary and Outlook

- ◆ The CLAS Collaboration at JLab has a comprehensive physics program to search for evidence of pentaquark states in photon induced reactions
- ◆ g11 ran in May–July 2004 accumulating a total integrated luminosity of  $70 \text{ pb}^{-1}$  in the photon energy range 1.6–3.8 GeV
- ◆ Preliminary results for the reaction  $\gamma p \rightarrow \Theta^+ \bar{K}^0$  show no indication of a narrow resonance in this channel
- ◆ An upper limit of 1–4 nb on the total cross section for this reaction was derived in contrast to the SAPHIR results
- ◆ Analysis of other reactions searching for the  $\Theta^+$  and  $\Theta^{++}$  are in progress
- ◆ A second experiment at higher energy designed to verify our previously published results in the reaction  $\gamma p \rightarrow \Theta^+ K^- \pi^+$  is planned